

الحلول

نعلم أن

I

$$3,16 < \sqrt{10} < 3,17$$

لدينا :

$$10^2 \times 3,16 < 10^2 \sqrt{10} < 10^2 \times 3,17$$

$$316 < \sqrt{10^5} < 317$$

$$316^2 < 10^5 < 317^2$$

إذن الأعداد الصحيحة الطبيعية التي مربعاتها أصغر من 10^5 هي 0 و 1 و 2 و

و 315 و 316 أي 317 عددا

$$\begin{aligned}
 S &= \frac{5!}{\sqrt{123} - \sqrt{3}} - \frac{-1}{\sqrt{3!} + \sqrt{7}} + \sqrt{6} \cdot \sqrt{7} \cdot \sqrt{123} \\
 &= \frac{120}{\sqrt{123} - \sqrt{3}} + \frac{1}{\sqrt{7} + \sqrt{6}} + \sqrt{6} \cdot \sqrt{7} \cdot \sqrt{123} \\
 &= \frac{120(\sqrt{123} + \sqrt{3})}{123 - 3} + \frac{\sqrt{7} \cdot \sqrt{6}}{7 - 6} + \sqrt{6} \cdot \sqrt{7} \cdot \sqrt{123} \\
 &= \sqrt{123} + \sqrt{3} + \sqrt{7} \cdot \sqrt{6} + \sqrt{6} \cdot \sqrt{7} \cdot \sqrt{123} \\
 &= \boxed{\sqrt{3}}
 \end{aligned}$$

$$\begin{array}{ll}
 \begin{array}{l}
 x < y \\
 x^2 < y^2 \\
 bx^2 < by^2 \\
 bx^2 + axy < by^2 + axy \\
 x(bx + ay) < y(by + ax)
 \end{array} &
 \begin{array}{l}
 x < y \\
 x^2 < y^2 \\
 ax^2 < ay^2 \\
 ax^2 + bxy < ay^2 + bxy \\
 x(ax + by) < y(ay + bx)
 \end{array}
 \end{array}$$

III

(2)

$$\boxed{\frac{x}{y} < \frac{ax+by}{bx+ay}}$$

(1)

$$\boxed{\frac{ax+by}{bx+ay} < \frac{y}{x}}$$

$$\boxed{\frac{x}{y} < \frac{ax+by}{bx+ay} < \frac{y}{x}}$$

من (1) و (2) نستنتج أن :


IV

$$\begin{aligned}
 P &= \left(1 - \frac{1}{4}\right) \left(1 - \frac{1}{9}\right) \left(1 - \frac{1}{16}\right) \dots \left(1 - \frac{1}{625}\right) \\
 &= \left(1 - \frac{1}{2}\right) \left(1 + \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 + \frac{1}{3}\right) \left(1 - \frac{1}{4}\right) \left(1 + \frac{1}{4}\right) \dots \left(1 - \frac{1}{25}\right) \left(1 + \frac{1}{25}\right) \\
 &= \left[\left(1 - \frac{1}{2}\right) \left(1 - \frac{1}{3}\right) \left(1 - \frac{1}{4}\right) \dots \left(1 - \frac{1}{25}\right)\right] \left[\left(1 + \frac{1}{2}\right) \left(1 + \frac{1}{3}\right) \left(1 + \frac{1}{4}\right) \dots \left(1 + \frac{1}{25}\right)\right] \\
 &= \left(\frac{1}{2} \times \frac{2}{3} \times \frac{3}{4} \times \dots \times \frac{24}{25}\right) \left(\frac{3}{2} \times \frac{4}{3} \times \frac{5}{4} \times \dots \times \frac{26}{25}\right) \\
 &= \frac{1}{25} \times \frac{26}{2} \\
 &= \frac{1}{25} \times 13 \\
 &= \boxed{\frac{13}{25}}
 \end{aligned}$$

لدينا :

$$\begin{aligned}
 AB &< OA + OB \\
 BC &< OB + OC \\
 CD &< OC + OD \\
 AD &< OA + OD
 \end{aligned}$$

V



$$\begin{aligned}
 p &< 2(OA + OC) + 2(OB + OD) \\
 p &< 2AC + 2BD \\
 p &< 2(AC + BD)
 \end{aligned}$$

(1) $\boxed{\frac{1}{2}p < AC + BD}$

$AC < AB + BC$: $\triangle ABC$ في المثلث

$AC < AD + CD$: $\triangle ACD$ في المثلث

$$2AC < AB + BC + AD + CD$$

$$2AC < p$$

$$\boxed{AC < \frac{1}{2}p}$$

$$\boxed{BD < \frac{1}{2}p}$$

نبرهن بنفس الطريقة أن

بما أن $AC + BD < \frac{1}{2}p + \frac{1}{2}p$ فإن $BD < \frac{1}{2}p$ و $AC < \frac{1}{2}p$

(2) $\boxed{AC + BD < p}$

$$\boxed{\frac{1}{2}p < AC + BD < p}$$

من (1) و (2) نستنتج أن :